

POSTOPERATIVE INSTRUCTIONS SHOULDER ARTHROPLASTY

WOUND CARE

- Keep your waterproof operative dressing clean and dry at all times
- If your dressing becomes wet or soiled, remove it and place a regular bandaid or gauze over incision
- As long as your dressing is intact, you may shower 48 hours after surgery
- DO NOT submerge incisions in water (bath, pool, hot tubs, etc) for at least 6 weeks after surgery
- DO NOT apply any cream/oil/lotion etc on or near your incision at any time
- It is normal for the shoulder to have some bleeding and swelling following surgery
- It is normal for your hand and fingers to swell and your arm to develop a bruise/discoloration

SLING USE

- You can come out of the sling if you are just relaxing on the couch, etc. You may also come out of the sling for showering, hygiene purposes and gentle activities (pendulums, elbow flexion/extension, wrist flexion/extension, opening and closing fist)
- The sling should always be worn in public and to sleep

ACTIVITY

- No driving until instructed otherwise by physician, it is illegal to drive in a sling
- May return to sedentary work 3-4 days after surgery, if pain is tolerable
- You are to **avoid** any PUSHING, PULLING, LIFTING greater than the weight of a cup of coffee
- Some patients have an easier time sleeping in a chair or recliner initially after surgery

MEDICATIONS

- Please follow the attached medication tracking sheet provided
- Most patients will require some narcotic pain medication for a short period of time
- You are also given prescriptions for extra-strength Tylenol and an anti-inflammatory (Celebrex). Use these first to reduce the amount of narcotic medication needed
- In most cases you will also be provided with prescriptions for Aspirin (to prevent blood clots), Omeprazole (to protect stomach while taking NSAIDs), Zofran (if needed for nausea), Colace (recommended to prevent constipation), and a Vitamin D/Calcium supplement (to enhance healing)
- Common side effects of the pain medication are nausea, drowsiness, and constipation – to decrease the side effects, take medication with food
- DO NOT drive or operate heavy machinery while taking narcotic medication. Do not consume alcohol or other recreational drugs while taking narcotic medication.

ICE THERAPY

- Use an icing machine or ice packs for 20 minutes every hour to help reduce pain and swelling

EXERCISE

- Begin exercises (pendulums and active flexion/extension at the elbow, wrist and fingers) 24 hours after surgery. Perform these 2-3 times per day
- Formal physical therapy (PT) typically begins after you are seen at your first post-operative appointment. A prescription and protocol will be provided at that time if not before

PAIN MANAGEMENT STRATEGIES

- Getting adequate rest, using your sling and icing your shoulder are essential
- Using all medications as prescribed will provide ‘multi-modal’ pain control pathways
- CBD (without THC) is helpful for some patients (oral intake only, DO NOT apply topically)
- Meditation is proven to reduce surgical pain (Headspace app has many relevant sessions)

FOLLOW-UP CARE/QUESTIONS


- Your first post operative visit will be 7-14 days after surgery
- Dr. Reynolds’ office can be reached at 336-333-6443, M-F, 8am – 4pm. MyWakeHealth messages will also be answered during these hours
- Outside of the above hours, you may call the same phone number if you have an emergency to reach the physician on call via an answering service

EMERGENCIES

- Contact Dr. Reynolds’ office at 336-333-6443 if any of the following are present:
 - Painful swelling or numbness (note that some swelling and numbness is normal)
 - Unrelenting pain
 - Fever (over 101° it is normal to have a low-grade fever the first day or two after surgery)
 - Redness around incisions
 - Color change in arm and/or hand
 - Continuous drainage or bleeding from incision (a small amount of drainage is expected)
 - Difficulty breathing
 - Excessive nausea/vomiting

If you have an emergency that requires immediate attention, proceed to the emergency room

Pendulum Exercises:



- While holding onto a table or the back of a chair with your good arm, bend forward and let your injured arm hang straight down
- This uses your legs and your hips to create movement that makes your arm swing freely
- Using the momentum from your hips and legs, guide the slightly swinging arm back and forth like a pendulum (or elephant trunk), then in circles that start small (about the size of a dinner plate) and gradually grow larger each day as pain allows